

CHAPTER # 1 NUMBER SYSTEM

1. $\sqrt{3}$ is a _____ number.


- (a) Odd number (b) Complex number (c) Rational (d) irrational

2. The multiplicative inverse of complex number $(0, 1)$ is

- (a) $(0, -1)$ (b) $(-1, 0)$ (c) $(1, 0)$ (d) $(1, 1)$

3. $(-i)^{19} = \dots\dots\dots$

- (a) $-i$ (b) i (c) 1 (d) -1

4. $a > 0$ 

- (a) $-a > 0$ (b) $-a < 0$ (c) $1/a > 0$ (d) $1/a < 0$

5. Modulus of complex number $3 - 4i$ is

- (a) 4 (b) 5 (c) -5 (d) 0

Answers:

- 1 D
2 A
3 B
4 C
5 B

CHAPTER # 2 SETS FUNCTIONS AND GROUPS

1. The tabular form of $\{x | x \in Q, x^2 = 3\}$ is

- (a) $\{9\}$ (b) $\{\sqrt{3}\}$ (c) $\{\sqrt{3}, -\sqrt{3}\}$ (d) $\{\}$

2. $A \cap B = \phi$, then $n(A-B)$ is equal to

- (a) $n(A)$ (b) $n(A \cap B)$ (c) $n(B)$ (d) $n(A \cup B)$

3. The power set of the empty set is

- (a) Empty set (b) non empty set (c) Power set (d) Improper set

4. $A \cup (A \cap B)$

- (a) B (b) $A \cap B$ (c) ϕ (d) A

5. The conjunction of two statements p and q is denoted by

- (a) $p \leftrightarrow q$ (b) $p \rightarrow q$ (c) $p \leftarrow q$ (d) $p \wedge q$

6. $p \wedge q$ is true if both are true. This is known as :

(a) Conjunction (b) negation (c) disjunction (d) none of these

7. Inverse of $p \rightarrow q$ is

(a) $\sim p \rightarrow \sim q$ (b) $p \rightarrow \sim q$ (c) $p \leftarrow q$ (d) $\sim q \rightarrow \sim p$

8. The converse of contrapositive of $p \rightarrow q$ is

(a) $q \rightarrow p$ (b) $\sim q \rightarrow \sim p$ (c) $\sim p \rightarrow \sim q$ (d) $p \rightarrow q$

9. A function which is onto is called

(a) Objective (b) Injective (c) bijective (d) surjective

10. Number of identity elements in any group is

(a) 1 (b) 2 (c) 3 (d) None of these

11. The set N w.r.t addition is a

(a) Group (b) monoid (c) null set (d) semi group

Answers:

1 D

2 A

3 B

4 D

5 D

6 D

7 A

8 A

9 C

10 A

11 D

CHAPTER # 3 MATRICES AND DETERMINANTS

1. For non-trivial solution $|A|$ is

(a) $|A| > 0$ (b) $|A| < 0$ (c) $|A| = 0$ (d) None

2. Minimum number of equation for any system of equations

(a) 2 (b) 3 (c) 4 (d) 10

3. The square matrix A is skew-Hermitian when $(A)^t = \dots\dots\dots$

(a) A (b) $-A$ (c) A (d) $-A$

4. The matrix $B = \begin{bmatrix} 1 & 4 \\ 4 & 8 \end{bmatrix}$ -----

(a) Singular matrix (b) Non Singular matrix (c) Symmetric matrix (d) None of these

6) If ω, ω^2 are complex cube roots of unity Then $\omega + \omega^2 =$

- A) 1 B) -1
C) 0 D) none of these

7) The roots of quadratic equation $x^2 - 4x = 0$ are

- A) Imaginary B) Rational & Different
C) Irrational D) Rational & Equal

8) If the area of a rectangle is 56 & the length is one more than the breadth then the dimensions are

- A) -8, -7 B) 8, 7
C) 14, 4 D) 28, 2

9) If one root of $4x^2 + 7hx - h^2 + 9 = 0$ is zero then $h =$

- A) 0 B) 3
C) -3 D) ± 3

10) The value of ω^{12} is

- A) 1 B) ω
C) ω^2 D) 0

Answers:

1. B
2. D
3. A
4. D
5. D
6. B
7. B
8. B
9. D
10. A

CHAPTER 5: PARTIAL FRACTIONS

1. An open formed by using the sign of equality “=” is called _____

- a) Equation
- b) In – equation
- c) True sentence
- d) False sentence

2. $2x = 3$ is a conditional equation it is true for _____

- a) 2
- b) 3
- c) $2/3$
- d) $3/2$

3. $x^2 + x - 6 = 0$ is a conditional equation and it is true for

- a) 2, 3
- b) 2, - 3
- c) - 2, - 3
- d) - 2, 3

4. The symbol _____ shall be used both for equation and identity

- a) \cong
- b) =
- c) \neq
- d) \equiv

5. $\frac{P(x)}{Q(x)}$, $Q(x) \neq 0$ is known as

- a) improper rational fraction
- b) rational fraction
- c) proper rational fraction
- d) none of the above

6. $\frac{9x^2}{x^3-1}$ is a fraction.

- a) rational fraction
- b) improper fraction
- c) rational fraction
- d) none of these

7. $\frac{x^2-3}{3x+1}$ is a fraction

- a) rational fraction
- b) proper fraction
- c) improper rational fraction
- d) none of these

8. There are _____ types of rational fraction .

- a) three
- b) four
- c) five
- d) two

9. The partial fraction of $\frac{1}{x^2-1}$ is

- a) $\frac{1}{2(x-1)} - \frac{1}{2(x+1)}$
- b) $\frac{1}{2(x-1)}$
- c) $\frac{1}{2(x+1)}$
- d) $\frac{1}{2(x-1)} + \frac{1}{2(x+1)}$

10. The partial fraction of $\frac{2x^2-3x+4}{(x-3)^3}$ is

- a) $\frac{2}{x-1}$
- b) $\frac{1}{(x-1)^2}$
- c) $\frac{2}{x-1} + \frac{1}{(x-1)^2} + \frac{3}{(x-1)^3}$
- d) $\frac{3}{(x-1)^3}$

Answers:

1. A
2. C
3. A
4. B
5. B
6. B
7. C
8. D
9. A
10. C

CHAPTER: 6 SEQUENCE AND SERIES

1. Which of the following is an A.P?

- (a) 2, 4, 6, (b) 1, 1/2, 1/3, (c) 1, 5, 11, 18, (d) 1, 4, 9,

2. The general term of the sequenced 2, 4, 6, 8, is

- (a) n (b) $2n$ (c) $2n - 1$ (d) n^2

3. The 8th term of the sequenced 5, 9, 13, is

- (a) 36 (b) 30 (c) 33 (d) 27

4. Which term of the sequenced 3, 9, 15, is 117

- (a) 19 (b) 20 (c) 21 (d) 22

5. The sum of the series $-1+1+3+\dots+21$ is _____

- (a) 110 (b) 120 (c) 130 (d) 140

6. The A. M b/w a & b is

- (a) $(a + b) / 2$ (b) $2ab / (a + b)$ (c) $(a - b) / 2$ (d) $\pm \sqrt{ab}$

7. The G.M b/w a & b is _____

- (a) $2ab / a + b$ (b) $(a + b) / 2$ (c) $\pm \sqrt{ab}$ (d) None

8. G. M b/w 2 & 8 is _____

- (a) 5 (b) ± 3.2 (c) - 4 (d) ± 4

9. A. M b/w 4 & 8 is..... _____

- (a) $\sqrt{32}$ (b) $-\sqrt{32}$ (c) 6 (d) $64 / 16$

10. If A, G & H are A.M, G.M & H.M b/w two numbers respectively then

- (a) $A < G < H$ (b) $A > G < H$ (c) $A < G > H$ (d) $A > G > H$

11. A geometric series cannot contain _____ as a term.

- (a) -1 (b) 1 (c) 0 (d) None

12. The sequence -1, 1, -1, 1.....is _____

- (a) A.P (b) G.P (c) H.P (d) None

13. The arrangement of number formed according to some definite rule is called _____

- (a) Sequence (b) Function (c) Series (d) None

Answers:

- 1. a
- 2. b
- 3. c
- 4. b
- 5. b
- 6. a
- 7. c
- 8. d
- 9. c
- 10. d
- 11. c
- 12. b
- 13. a

CHAPTER 7: PERMUTATIONS, COMBINATIONS AND PROBABILITY

1. For a positive integer n $n!$ = _____

- (a) n^2 (b) $n(n-1)(n-2).....3.2.1$ (c) $n.n(n-1)...2.1$ (d) $n.5.2.1$

2. $0!$ =

- (a) 1 (b) 0 (c) undefined (d) None

3. ${}^n P_0$ =

- (a) 1 (b) 0 (c) 0 (d) $n!$

4. ${}^n P_r$ =

(a) $n(n-1)\dots(n-r-1)$ (b) $n(n-1)(n-2)\dots(n-r+1)$ (c) $n(n-1)\dots(n-r)$ (d) None

5. ${}^{20}P_3 = \underline{\hspace{2cm}}$

(a) $20 \times 19 \times 18$ (b) 20×3 (c) $20! / 3!$ (d) $3 / 20!$

6. ${}^nP_{n-1} = \underline{\hspace{2cm}}$

(a) $(n-1)!$ (b) $n(n-1)$ (c) $n!$ (d) 1

7. If ${}^nP_2 = 30$ then $n =$

(a) 5 (b) 6 (c) 2 (d) 28

8. In how many different ways can eight books be arranged on a shelf?

(a) $8!$ (b) 8 (c) $56!$ (d) 56

9. The number of committees of seven person formed from a group of 10 persons will be

(a) 71 (b) 101 (c) 100 (d) 120

10. If A & B are mutually disjoint events then $P(A \cup B) =$

(a) $P(A) + P(B) - P(A \cap B)$ (b) $P(A) - P(B)$ (c) $P(A) + P(B)$ (d) None

Answers:

1. b
2. a
3. a
4. b
5. a
6. c
7. b
8. a
9. d
10. c

CHAPTER 8: MATHEMATICAL INDUCTION AND BINOMIAL THEOREM

1) If x is so small that its square and higher powers be neglected then $(1 + 3x)^{-2} =$

- A) $1 + 9x$ B) $1 - 9x$
C) $1 + 6x$ D) $1 - 6x$

2) For every positive integer n $1 + 5 + 9 + \dots + (4n - 3)$ is equal to

- A) $n(2n - 1)$ B) $(2n - 1)$
D) $n - 1$ D) n

3) The number of terms in the expansion of $(2x + y)^6$ are

- A) 6 B) 7
C) 8 D) 14

4) The term involving x^4 in the expansion of $(3 - 2x)^7$ is

- A) 120 B) 1512
C) 1250 D) 15120

5) $(a + x)^n = \sum_{r=0}^n \binom{n}{r} a^{n-r}$ where a and x are:

- A) imaginary B) Rational
C) Irrational D) Real numbers

6) The expansion $(1 + x)^{-3}$ holds when

- A) $|x| > 1$ B) $|x| < 1$
C) $|x| > 1$ D) $x < 1$

7) $1 + 2 + 3 + \dots + n =$

- A) $\frac{n^2(n+1)^2}{4}$ B) $\frac{n(n+1)}{2}$
C) $\frac{n(n+1)(2n+1)}{6}$ D) $\frac{n^2}{2}$

8) Number of terms in the expansion of $(a + x)^n$ is

- A) $n - 1$ B) $n + 1$
C) $n + 2$ D) $n + 3$

9) The middle term of the expansion $(1 + 2x)^6$ is

A) 1st term

B) 4th term

C) 2nd term

D) 3rd term

10) If n is added to the expansion $(a + x)^n$ has middle terms.

A) 2

B) 3

C) 4

D) 5

Answers:

1. D

2. A

3. A

4. D

5. D

6. B

7. B

8. A

9. B

10. A

CHAPTER 9: FUNDAMENTALS OF TRIGONOMETRY

1. What is the length of an arc of a circle of radius 5cm, whose central angle is of 140°

a) 2.443 radians

b) 1.443 radians

c) 0.443 radians

d) 2 radians

e) None of these

2. If a circle is divided into 360 parts, then the angle subtended by each part at the center of the circle is called _____

a) 1 radian

b) 1 degree

c) 1 angstrom

d) 1 minute

e) None of these

3. The union of two non-collinear rays which have a common endpoint is called the

a) Angle

b) Radian

c) Degree

d) Minute

e) Second

4. The 60th part of one degree is called one

- a) centimeter
- b) radian
- c) degree
- d) minute
- e) none of these

5. $1^\circ =$ _____

- a) 60'
- b) 60''
- c) 3600'
- d) 360'
- e) None of these

6. The system of measurement in which the angle is measured in degrees, and its subunits, minutes and seconds is called

- a) Circular system
- b) Sexagesimal system
- c) MKS system
- d) CGS system
- e) None of these

7. The length of the arc cut off on a circle of radius 6cm by a central angle of $2\pi/3$ radians

- a) 12.566cm
- b) 10.033cm
- c) 12.113cm
- d) 9.156cm
- e) 6.56cm

8. The radius of the circle when $l = 3\text{cm}$, $\theta = 3.4$ radians.

- a) 0.214 cm
- b) 9.419 cm
- c) 3.146 cm
- d) 4.978 cm
- e) None of these

9. $\sin\theta = \frac{\text{Perpendicular}}{?}$

- a) base
- b) hypotenuse
- c) $\cos\theta$
- d) $\tan\theta$
- e) none of these

10. The pendulum of a clock is 40cm long and it swings through an angle of 30° each second. How far does the tip of the pendulum move in 1 seconds?

- a) 10cm
- b) 15.71cm
- c) 20.94cm
- d) 28.65cm
- e) 40cm

Answers:

- 1. E
- 2. B
- 3. A
- 4. D
- 5. A
- 6. B
- 7. A
- 8. E
- 9. B
- 10. C

CHAPTER 10: TRIGNOMETRIC IDENTITIES

1. $\sin(\alpha - \beta) = \dots\dots\dots$

- (a) $\sin \alpha \cos \beta + \cos \alpha \sin \beta$
- (b) $\sin \alpha \cos \beta - \cos \alpha \sin \beta$
- (c) $\cos \alpha \cos \beta - \sin \alpha \sin \beta$
- (d) $\cos \alpha \cos \beta + \sin \alpha \sin \beta$

2. $\cos(\alpha + \beta) = \dots\dots\dots$

- (a) $\sin \alpha \cos \beta + \cos \alpha \sin \beta$
- (b) $\sin \alpha \cos \beta - \cos \alpha \sin \beta$
- (c) $\cos \alpha \cos \beta - \sin \alpha \sin \beta$
- (d) $\cos \alpha \cos \beta + \sin \alpha \sin \beta$

3. $\tan(-\beta) = \dots\dots\dots$

- (a) $\tan \beta$
- (b) $-\tan \beta$
- (c) $\cot \beta$
- (d) $\cos \beta$

4. $\cos(2700 - \theta) = \dots\dots\dots$

- (a) $-\sin \theta$ (b) $\sin \theta$ (c) $-\cos \theta$ (d) $\cos \theta$

5. $\tan(2700 + \theta) = \dots\dots\dots$

- (a) $\tan \theta$ (b) $-\tan \theta$ (c) $\cot \theta$ (d) $-\cot \theta$

6. $\sin(1800 - \theta) = \dots\dots\dots$

- (a) $\cos \theta$ (b) $\sin \theta$ (c) $-\sin \theta$ (d) $-\cos \theta$

7. $\tan(450 - \theta) = \dots\dots\dots$

- (a) $1 + \tan \theta$ (b) $\cos \theta + \sin \theta$ (b) $1 - \tan \theta \cos \theta - \sin \theta$
 (c) $1 - \tan \theta$ (d) $\sin \theta + \cos \theta$ (d) $1 + \tan \theta \sin \theta - \cos \theta$

8. $\cot 150 = \dots\dots\dots$

- (a) $\sqrt{3} - 1 / \sqrt{3} + 1$ (b) $1 - \sqrt{3} / 1 + \sqrt{3}$ (c) $1 + \sqrt{3} / 1 - \sqrt{3}$ (d) $\sqrt{3} + 1 / \sqrt{3} - 1$

9. $\cos(2000 40') = \dots\dots\dots$

- (a) $-\sin(200 40')$ (b) $\sin(200 40')$ (c) $\cos(200 40')$ (d) $-\cos 200 40'$

10. $\sin \alpha + \cos \alpha = \dots\dots\dots \sec 4\alpha \quad \operatorname{cosec} 4\alpha$

- (a) $\sec 5\alpha$ (b) $\sin 5\alpha$ (c) $\operatorname{cosec} 5\alpha$ (d) $\cos 5\alpha$

11. $\sin \beta = \dots\dots\dots$

- (a) $2\sin \beta \cos \beta$ (b) $\sin \beta / 2 \cos \beta / 2$ (c) $2\sin \beta / 2 \cos \beta / 2$ (d) None

12. $\cos 2\alpha = \dots\dots\dots$

- (a) $2\cos 2\alpha + 1$ (b) $2\cos 2\alpha - 1$ (c) $1 + 2\sin 2\alpha$ (d) $2\sin 2\alpha - 1$

13. $\cos^2 150 - \sin^2 150 = \dots\dots\dots$

- (a) $\cos^2 150$ (b) $\cos^4 300$ (c) $\frac{1}{2}$ (d) $\sqrt{3} / 2$

Answers:

1. a
2. c
3. b
4. a
5. d
6. b
7. c

- 8. d
- 9. d
- 10. b
- 11. c
- 12. b
- 13. d

CHAPTER 11: TRIGONOMETRIC FUNCTIONS AND THEIR GRAPHS

1. 'Sine' and 'cosine' are periodic function whose period is
 (a) $\pi/2$ (b) π (c) 2π (d) 4π
2. Tangent and cotangent are periodic function whose period is.....
 (a) $\pi/2$ (b) π (c) 2π (d) 4π
3. The period of sec and cosec function is
 (a) $\pi/2$ (b) $\pi/4$ (c) $3\pi/4$ (d) 2π
4. The domain of sine & cosine is
 (a) $-\pi/2 \leq x \leq \pi/2$ (b) $-\pi/2 < x < \pi/2$ (c) $0 \leq x \leq \pi$ (d) None
5. The range of $y = \sin x$ & $y = \cos x$ is
 (a) $0 \leq y \leq 1$ (b) $-1 \leq y \leq 0$ (c) $-1 \leq y \leq 1$ (d) $-1 < y < 1$
6. $\tan 180^\circ =$
 (a) 0 (b) 1 (c) -1 (d) ∞
7. $\sin(315^\circ) =$
 (a) $1/\sqrt{2}$ (b) $-1/\sqrt{2}$ (c) $\sqrt{3}/2$ (d) $-\sqrt{3}/2$
8. In 1st Quadrant the value of sine is.....
 (a) Increase from 0 to 1 (b) Increase from -1 to 0 (c) Decrease from 1 to 0 (d) Decrease from 0 to -1
9. In 3rd quadrant the value of cosines
 (a) Increases (b) decreases (c) Constant (d) None
10. $\cos(240^\circ) =$...
 (a) $\sqrt{3}$ (b) $-\sqrt{3}$ (c) $\frac{1}{2}$ (d) $-\frac{1}{2}$

Answers:

1. c
2. a
3. d
4. d
5. c
6. a

- 7. b
- 8. a
- 9. a
- 10. b

CHAPTER12: APPLICATION OF TRIGONOMETRY

1. How many important elements a triangle has.....

- A. 5
- B. 6
- C. -5
- D. 4
- E. None of these

2. Angle above the eye level

- a) Angle of elevation
- b) Angle of depression
- c) Constant angle
- d) Right angle
- e) Obtuse angle

3. A tree of 8m high has the shadow 6m in length, the angle of elevation of the sun at that moment is

- a) 0
- b) $53^{\circ}7'$
- c) 90°
- d) 180°
- e) 225°

4. At the top of a cliff 80m high, the angle of depression of a boat is 12° . the distance of the boat from the cliff is

- a) 100m
- b) 255m
- c) 377m
- d) 477m
- e) 733m

5. The area of a triangle with $a = 300$, $b=120$, $\gamma = 150^{\circ}$ is

- a) 5000 square units
- b) 6000 square units
- c) 7000 square units
- d) 9000 square units

6. a circle drawn inside a triangle and touching its sides is called the

- a) Circum circle
- b) In circle
- c) Escribed circle
- d) Normal
- e) None of these

7. The circle passing through three vertices of a triangle is called a

- a) Circum circle
- b) In circle
- c) Escribed circle
- d) Tangent
- e) None of these

8. $r_1 r_2 r_3 =$

- a) Rr^2
- b) rR^2
- c) Rs^2
- d) rR^2
- e) rs^2

9. $r_1 r_2 + r_2 r_3 + r_3 r_1 =$

- a) r_1^2
- b) Δ^2
- c) R^2
- d) r^2
- e) s^2

10. if Δ is the area of a triangle ABC, then $\Delta =$

- a) $\sqrt{s(s+a)(s-b)(s-c)}$
- b) $\sqrt{s(s-a)(s+b)(s-c)}$
- c) $\sqrt{s(s-a)(s-b)(s+c)}$
- d) $\sqrt{s(s+a)(s+b)s+c}$
- e) $\sqrt{s(s-a)(s-b)s-c}$

Answers:

- 1. B
- 2. A
- 3. B
- 4. C
- 5. D
- 6. B
- 7. A
- 8. E
- 9. E
- 10. E

CHAPTER 13: INVERSE TRIGONOMETRIC FUNCTIONS

- The inverse exists only for the function which is
(a) One to one (b) onto (c) into (d) All of these
- To make a trigonometric function one to one, it's _____ is restricted.
(a) Domain (b) period (c) Range (d) None
- The domain of $y = \sin^{-1}$ is.
(a) $(-\pi, \pi)$ (b) $(-\pi/2, \pi/2)$ (c) $(-1, 2)$ (d) None of these
- Inverse sine function is written as..... (
a) $(\sin x)^{-1}$ (b) $\sin 1$ (c) $\arcsin^{-1} x$ (d) \arcsin^{-1}
- The domain of $y = \sin^{-1} x$ function is...
(a) $(-1, 1)$ (b) $(-\infty, \infty)$ (c) $(0, \pi)$ (d) $(-\pi/2, \pi/2)$
- The range of $y = \sin^{-1}$ function is....
(a) $(-1, 1)$ (b) $(-\infty, \infty)$ (c) $(0, \pi)$ (d) $(-\pi/2, \pi/2)$
- The inverse cosine function can also be written as
(a) $(\cos x)^{-1}$ (b) $\cos x^{-1}$ (c) $\arcsin x$ (d) $\arcsin^{-1} x$
- The domain of $y = (\cos)^{-1} x$ function is...
(a) $(-1, 1)$ (b) $(-\infty, \infty)$ (c) $(0, \pi)$ (d) $(-\pi/2, \pi/2)$
- The range of $y = (\cos)^{-1} x$ function is...
(a) $(-1, 1)$ (b) $(-\infty, \infty)$ (c) $(0, \pi)$ (d) $(-\pi/2, \pi/2)$
- Inverse tangent function can be written also written as
(a) $(\tan x)^{-1}$ (b) $\tan x^{-1}$ (c) $\arcsin x$ (d) $\arcsin^{-1} x$

Answers:

- a
- a
- b
- c
- a
- d
- c
- a
- c
- c
- d
- a
- b

- 14. a
- 15. b
- 16. c
- 17. b
- 18. b
- 19. b
- 20. a
- 21. b

CHAPTER 14: SOLUTIONS OF TRIGONOMETRIC EQUATIONS

1. If $\sin x = \cos x$, then the general solution is

- (a) $n\pi$, where $n \in \mathbb{Z}$
- (b) $2n\pi$, where $n \in \mathbb{Z}$
- (c) $\pi/4 + 2n\pi$, where $n \in \mathbb{Z}$
- (d) $\pi/4$

2. Solution of the equation $\sin x = \sqrt{3}/2$, $x \in (0, \pi)$ are

- (a) $\pi/3, 2\pi/3$
- (b) $\pi/6, \pi/3$
- (c) $\pi/3, 5\pi/3$
- (d) None of these

3. Solution of the equation $1 + \cos x = 0$ is

- (a) $\pi/2$
- (b) π
- (c) 2π
- (d) None of these

4. The solution set of $\tan 2x = 1$ in $(0, \pi)$ is:

- (a) $\{\pi/8, 5\pi/8\}$
- (b) $\{\pi/4, 5\pi/4\}$
- (c) $\{\pi/4, 3\pi/4\}$
- (d) None of these

5. The solution set of $\sec x = -2$ in $(0, \pi)$ is:

- (a) $2\pi/3, 4\pi/3$
- (b) $\pi/3, 2\pi/3$
- (c) $\pi/3, 4\pi/3$
- (d) None of these

6. The solution set of $\sin x = 0$ is :

- (a) $2n\pi, n \in \mathbb{Z}$
- (b) $n\pi, n \in \mathbb{Z}$
- (c) $2n\pi, n \in \mathbb{Q}$
- (d) None of these

7. The solution of $\tan^2 x = 3$, which lie in $(0, \pi)$ is

(a) $4\pi/3, 5\pi/3$

(b) $\pi/3, 2\pi/3$

(c) $\pi/3, 4\pi/3$

(d) None of these

8. If $\tan(x/2) = \sqrt{3}$, and x lies in $(0, 2\pi)$, then x is equal to

(a) 0 and x

(b) $\pi/3$ and $5\pi/3$

(c) $\pi/6$ and $5\pi/6$

(d) None of these

9. General solution of the equation $1 + \cos x = 0$ is

(a) $\{\pi/2 + 2n\pi\}$

(b) $\{-\pi/2 + 2n\pi\}$

(c) $\{\pi + 2n\pi\}$

(d) None of these

10. $\cos x = 1/2$ has solution:

(a) $\pi/2$

(b) $\pi/3$

(c) $\pi/4$

(d) $\pi/6$

Answers:

1. C

2. A

3. B

4. A

5. A

6. B

7. D

8. B

9. C

10. D